


1. Energy is released when nitrogen gas combines with hydrogen gas to form ammonia according to the equation:




Which of the following is responsible for the release of energy?

- A. Breaking nitrogen bonds
 - B. Breaking hydrogen bonds
 - C. Forming nitrogen-hydrogen bonds
 - D. Breaking both nitrogen and hydrogen bonds.
 - E. All of these release energy
2. The standard enthalpy of formation of atomic oxygen is:
- A. Negative, since all spontaneous reactions are exothermic
 - B. Zero, since oxygen is an element
 - C. Positive, since breaking of bonds is always endothermic
 - D. Negative, zero, or positive depending on the temperature
 - E. Zero, since oxygen is an element and Positive, since breaking of bonds is always endothermic
3. Given the following standard enthalpies of formation:

Compound	ΔH_f (kJ/mol)
$\text{CH}_4(\text{g})$	-75
$\text{CO}_2(\text{g})$	-394
$\text{H}_2\text{O}(\text{g})$	-242


What is the standard enthalpy change for the combustion of methane:




- A. +561 kJ
- B. -561 kJ
- C. +803 kJ
- D. -803 kJ
- E. -711 kJ

4. The standard enthalpy of formation for $\text{H}_2\text{O}(\text{l})$ is $\Delta H = -285 \text{ kJ/mol}$.

Beginning with $\text{H}_2\text{O}(\text{l})$, which of the following physical and/or chemical changes is/are **exothermic**?



- A. only i
 B. i, ii & iii
 C. ii & iv
 D. only ii
 E. only iv
5. Which of the following are **NOT** correctly written as a standard enthalpy of formation reactions?




- A. i & ii
 B. iii & iv
 C. ii, iii & iv
 D. i & iv
 E. all are correct

6. 100 mL of water, 100 mL of alcohol, and 200 mL of an unknown liquid, all at 25°C, are heated at the **same rate under identical conditions**. After 3 minutes the temperature of the alcohol is 50°C. It took 5 minutes for the water to reach 50°C. It took 6 minutes for the unknown liquid to reach 50°C.

Which statement is TRUE regarding the specific heat, C_s ?

- A. C_s (water) = C_s (alcohol)
- B. C_s (water) > C_s (alcohol)
- C. C_s (water) < C_s (unknown)
- D. C_s (alcohol) > C_s (unknown)

7. What amount of heat energy, kJ, is released in the combustion of 16.0 g of C_3H_6 ?





- A. 581
- B. 783
- C. 504
- D. 624
- E. 725


8. Which of these will lead to the greatest increase in temperature when added to 100.0 mL of water at 25°C?

- A. 94 g of gold at 150°C, specific heat = 0.128 J/g K.
- B. 50 g of iron at 110°C, specific heat = 0.452 J/g K.
- C. 66 g of silver at 124°C, specific heat = 0.234 J/g K.
- D. 70 g of copper at 150°C, specific heat = 0.385 J/g K.

9. Given the following equations and ΔH° values, determine the heat of reaction (kJ) at 298 K for the reaction:



- A. +770.8
 B. -605.0
 C. -300.1
 D. +291.2
 E. -291.6
10. A volume of 50.0 mL of 0.400 M NaOH(aq) was added to 20.0 mL of 0.500 M $\text{H}_2\text{SO}_4(\text{aq})$ in a calorimeter of heat capacity 39.0 JK^{-1} . The temperature of the resulting solution rose by 3.60°C . What is the total heat produced (in kJ) in this neutralization? Assume that the specific heat capacity of the neutralized solution is $4.18 \text{ JK}^{-1}\text{g}^{-1}$.



- A. -102
 B. -111
 C. -119
 D. -126
 E. -132

11. In the introduction to Chapter 6 I played several clips from the hit TV show “The Big Bang Theory” in class. In this episode, Dr. Sheldon Cooper is trying to visualize the behavior of electrons traveling through a graphene sheet. Which of the following statements is/are true regarding his findings?

- i) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model with balls from a ball pit after he determined the proper size ratio with bigger carbon atoms.
- ii) Just like Dalton, Thompson, and Bohr, it was relatively easy for Sheldon to develop an accurate model of the behavior of an electron.
- iii) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model after recognizing the electron behaved as both a particle and a wave.

- A. ii only
- B. iii only
- C. i & ii
- D. i & iii
- E. ii & iii


12. What is the energy of the photons emitted from the He-Ne laser, wavelength = 655 nm?

- A. 2.27×10^{-19} J
- B. 4.34×10^{-40} J
- C. 3.03×10^{-19} J
- D. 1.01×10^{-27} J
- E. 5.29×10^{-18} J

13. Which emission line in the hydrogen spectrum occurs at highest frequency?

- A. $n = 3$ to $n = 1$
- B. $n = 1$ to $n = 3$
- C. $n = 4$ to $n = 2$
- D. $n = 2$ to $n = 4$
- E. $n = 5$ to $n = 7$

14.



Suppose a certain atom has four energy levels, and assume all transitions between levels are possible. Which statement is TRUE?

- A. The lowest energy light is emitted for $n=3 \rightarrow n = 1$
 - B. The longest wavelength light is emitted for $n = 4 \rightarrow n = 2$
 - C. The potential energy is greatest when the electron is at the $n=1$ level.
 - D. There are a total of 6 spectral emission lines.
 - E. There are a total of 4 spectral emission lines.
15. Sodium metal requires a photon with a minimum energy of 4.41×10^{-19} J to emit electrons. Suppose you have two lasers, one emitting light with a wavelength of 210 nm and the other emitting light with a wavelength of 335 nm. Each laser produces 10,000 photons/sec.

Which statement is true regarding the removal of electrons from the metal's surface?


- A. Both lasers eject electrons from the metal surface at the same rate.
- B. Both lasers eject electrons from the metal surface, but the weaker laser needs to operate for a longer period of time.
- C. Only the laser with photons of 335 nm ejects electrons.
- D. Only the laser with photons of 210 nm ejects electrons.
- E. Neither laser will eject electrons.

16. Which of the following statements best describes a $4d_{xz}$ atomic orbital?
- The $4d_{xz}$ orbital has the same number of radial and angular nodes.
 - The $4d_{xz}$ orbital has more radial nodes than angular nodes.
 - The $4d_{xz}$ orbital has more angular nodes than radial nodes.
 - The $4d_{xz}$ orbital does not have any radial nodes.
 - The $4d_{xz}$ orbital does not have any angular nodes.
17. Which of the following electron configurations is correct for mercury (Hg)?
- $[\text{Kr}] 6s^2 6f^{14} 6d^{10}$
 - $[\text{Xe}] 4s^2 5f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 5d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 5d^{10}$
18. How many electrons in a zinc atom have the quantum number $m_l = 0$?
- 6
 - 8
 - 10
 - 14
 - 30
19. Which response correctly fills in the blanks?
- The atom _____ has _____ core electrons, _____ valence electrons, and _____ unpaired electrons.
- oxygen, 2, 6, 4
 - chlorine, 12, 5, 2
 - nitrogen, 2, 5, 3
 - neon, 2, 8, 6
 - fluorine, 2, 7, 5
20. Which of the following sets of quantum numbers could describe one of the valence electrons in a sulfur atom in the ground state?
- $n=1 \quad l=0 \quad m_l=0 \quad m_s=+\frac{1}{2}$
 - $n=2 \quad l=1 \quad m_l=-1 \quad m_s=-\frac{1}{2}$
 - $n=3 \quad l=1 \quad m_l=0 \quad m_s=+\frac{1}{2}$
 - $n=3 \quad l=3 \quad m_l=-2 \quad m_s=+\frac{1}{2}$
 - $n=4 \quad l=0 \quad m_l=-1 \quad m_s=+\frac{1}{2}$

21. The attraction of the nucleus on the outermost electron in an atom tends to:
- decrease moving from left to right and top to bottom on the periodic table.
 - decrease moving from right to left and top to bottom on the periodic table.
 - decrease moving from left to right and bottom to top on the periodic table.
 - decrease moving from right to left and bottom to top of the periodic table.
 - the attraction of the nucleus to the outermost electron remains constant for each element on the periodic table.
22. Lithium's first and second ionization energies are 519 kJ/mol and 7300 kJ/mol, respectively. Element X has a first ionization energy of 590 kJ/mol and a second ionization energy of 1150 kJ/mol. Element X is most likely to be:
- Oxygen
 - Sodium
 - Calcium
 - Xenon
 - Fluorine
23. Which of the following is correctly ordered in terms of atomic radius, from smallest to largest?
- $\text{Mg}^{2+} < \text{Na}^+ < \text{F}^-$
 - $\text{Mg}^{2+} < \text{F}^- < \text{Na}^+$
 - $\text{F}^- < \text{Na}^+ < \text{Mg}^{2+}$
 - $\text{F}^- < \text{Mg}^{2+} < \text{Na}^+$
 - $\text{Na}^+ < \text{F}^- < \text{Mg}^{2+}$
24. Arrange the following in order of increasing electron affinity: Cl, P, S, Ar.
- Most positive EA $\text{Cl} < \text{S} < \text{P} < \text{Ar}$ Most negative EA
 - Most positive EA $\text{Ar} < \text{Cl} < \text{S} < \text{P}$ Most negative EA
 - Most positive EA $\text{P} < \text{S} < \text{Cl} < \text{Ar}$ Most negative EA
 - Most positive EA $\text{Ar} < \text{P} < \text{S} < \text{Cl}$ Most negative EA
 - Most positive EA $\text{Cl} < \text{Ar} < \text{P} < \text{S}$ Most negative EA
25. Which of the following statements is TRUE?
- The first ionization potential of H is greater than that of He.
 - The ionic radius of Fe^+ is larger than that of Fe^{3+} .
 - The electron affinity of C is greater than that of Cl.
 - The atomic radius of Li is larger than that of Cs.
 - All are false.

Version: B

1. Energy is released when nitrogen gas combines with hydrogen gas to form ammonia according to the equation:




Which of the following is responsible for the release of energy?

- A. Forming nitrogen-hydrogen bonds
 - B. All of these release energy
 - C. Breaking nitrogen bonds
 - D. Breaking both nitrogen and hydrogen bonds.
 - E. Breaking hydrogen bonds
2. The standard enthalpy of formation of atomic oxygen is:
- A. Positive, since breaking of bonds is always endothermic
 - B. Negative, zero, or positive depending on the temperature
 - C. Zero, since oxygen is an element and Positive, since breaking of bonds is always endothermic
 - D. Zero, since oxygen is an element
 - E. Negative, since all spontaneous reactions are exothermic
3. Given the following standard enthalpies of formation:

Compound	ΔH_f (kJ/mol)
CH ₄ (g)	-75
CO ₂ (g)	-394
H ₂ O(g)	-242


What is the standard enthalpy change for the combustion of methane:




- A. +803 kJ
- B. +561 kJ
- C. -561 kJ
- D. -803 kJ
- E. -711 kJ

4. The standard enthalpy of formation for $\text{H}_2\text{O}(\text{l})$ is $\Delta\text{H} = -285 \text{ kJ/mol}$.

Beginning with $\text{H}_2\text{O}(\text{l})$, which of the following physical and/or chemical changes is/are **exothermic**?



- A. ii & iv
 B. only i
 C. only iv
 D. only ii
 E. i, ii & iii
5. Which of the following are **NOT** correctly written as a standard enthalpy of formation reactions?




- A. ii, iii & iv
 B. iii & iv
 C. i & ii
 D. i & iv
 E. all are correct

6. 100 mL of water, 100 mL of alcohol, and 200 mL of an unknown liquid, all at 25°C, are heated at the **same rate under identical conditions**. After 3 minutes the temperature of the alcohol is 50°C. It took 5 minutes for the water to reach 50°C. It took 6 minutes for the unknown liquid to reach 50°C.

Which statement is TRUE regarding the specific heat, C_s ?

- A. C_s (alcohol) > C_s (unknown)
- B. C_s (water) < C_s (unknown)
- C. C_s (water) > C_s (alcohol)
- D. C_s (water) = C_s (alcohol)

7. What amount of heat energy, kJ, is released in the combustion of 16.0 g of C_3H_6 ?





- A. 783
- B. 624
- C. 725
- D. 504
- E. 581


8. Which of these will lead to the greatest increase in temperature when added to 100.0 mL of water at 25°C?

- A. 66 g of silver at 124°C, specific heat = 0.234 J/g K.
- B. 94 g of gold at 150°C, specific heat = 0.128 J/g K.
- C. 50 g of iron at 110°C, specific heat = 0.452 J/g K.
- D. 70 g of copper at 150°C, specific heat = 0.385 J/g K.

9. Given the following equations and ΔH° values, determine the heat of reaction (kJ) at 298 K for the reaction:



- A. -291.6
 B. -300.1
 C. +291.2
 D. -605.0
 E. +770.8
10. A volume of 50.0 mL of 0.400 M NaOH(aq) was added to 20.0 mL of 0.500 M H_2SO_4 (aq) in a calorimeter of heat capacity 39.0 JK^{-1} . The temperature of the resulting solution rose by 3.60°C . What is the total heat produced (in kJ) in this neutralization? Assume that the specific heat capacity of the neutralized solution is $4.18 \text{ JK}^{-1}\text{g}^{-1}$.



- A. -111
 B. -126
 C. -119
 D. -102
 E. -132

11. In the introduction to Chapter 6 I played several clips from the hit TV show “The Big Bang Theory” in class. In this episode, Dr. Sheldon Cooper is trying to visualize the behavior of electrons traveling through a graphene sheet. Which of the following statements is/are true regarding his findings?

- i) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model with balls from a ball pit after he determined the proper size ratio with bigger carbon atoms.
- ii) Just like Dalton, Thompson, and Bohr, it was relatively easy for Sheldon to develop an accurate model of the behavior of an electron.
- iii) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model after recognizing the electron behaved as both a particle and a wave.

- A. i & iii
- B. i & ii
- C. iii only
- D. ii only
- E. ii & iii


12. What is the energy of the photons emitted from the He-Ne laser, wavelength = 655 nm?

- A. 4.34×10^{-40} J
- B. 2.27×10^{-19} J
- C. 3.03×10^{-19} J
- D. 1.01×10^{-27} J
- E. 5.29×10^{-18} J

13. Which emission line in the hydrogen spectrum occurs at highest frequency?

- A. $n = 1$ to $n = 3$
- B. $n = 4$ to $n = 2$
- C. $n = 3$ to $n = 1$
- D. $n = 2$ to $n = 4$
- E. $n = 5$ to $n = 7$

14.



Suppose a certain atom has four energy levels, and assume all transitions between levels are possible. Which statement is TRUE?

- A. The longest wavelength light is emitted for $n = 4 \rightarrow n = 2$
 - B. The potential energy is greatest when the electron is at the $n=1$ level.
 - C. The lowest energy light is emitted for $n=3 \rightarrow n = 1$
 - D. There are a total of 6 spectral emission lines.
 - E. There are a total of 4 spectral emission lines.
15. Sodium metal requires a photon with a minimum energy of 4.41×10^{-19} J to emit electrons. Suppose you have two lasers, one emitting light with a wavelength of 210 nm and the other emitting light with a wavelength of 335 nm. Each laser produces 10,000 photons/sec.

Which statement is true regarding the removal of electrons from the metal's surface?


- A. Only the laser with photons of 335 nm ejects electrons.
- B. Only the laser with photons of 210 nm ejects electrons.
- C. Both lasers eject electrons from the metal surface, but the weaker laser needs to operate for a longer period of time.
- D. Both lasers eject electrons from the metal surface at the same rate.
- E. Neither laser will eject electrons.

16. Which of the following statements best describes a $4d_{xz}$ atomic orbital?
- The $4d_{xz}$ orbital has more radial nodes than angular nodes.
 - The $4d_{xz}$ orbital does not have any angular nodes.
 - The $4d_{xz}$ orbital has more angular nodes than radial nodes.
 - The $4d_{xz}$ orbital does not have any radial nodes.
 - The $4d_{xz}$ orbital has the same number of radial and angular nodes.
17. Which of the following electron configurations is correct for mercury (Hg)?
- $[\text{Xe}] 4s^2 5f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 5d^{10}$
 - $[\text{Kr}] 6s^2 6f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 5d^{10}$
18. How many electrons in a zinc atom have the quantum number $m_l = 0$?
- 10
 - 6
 - 30
 - 14
 - 8
19. Which response correctly fills in the blanks?
- The atom _____ has _____ core electrons, _____ valence electrons, and _____ unpaired electrons.
- nitrogen, 2, 5, 3
 - oxygen, 2, 6, 4
 - chlorine, 12, 5, 2
 - neon, 2, 8, 6
 - fluorine, 2, 7, 5
20. Which of the following sets of quantum numbers could describe one of the valence electrons in a sulfur atom in the ground state?
- $n=3 \quad l=1 \quad m_l=0 \quad m_s=+\frac{1}{2}$
 - $n=4 \quad l=0 \quad m_l=-1 \quad m_s=+\frac{1}{2}$
 - $n=2 \quad l=1 \quad m_l=-1 \quad m_s=-\frac{1}{2}$
 - $n=3 \quad l=3 \quad m_l=-2 \quad m_s=+\frac{1}{2}$
 - $n=1 \quad l=0 \quad m_l=0 \quad m_s=+\frac{1}{2}$

21. The attraction of the nucleus on the outermost electron in an atom tends to:
- decrease moving from left to right and bottom to top on the periodic table.
 - decrease moving from right to left and top to bottom on the periodic table.
 - decrease moving from left to right and top to bottom on the periodic table.
 - decrease moving from right to left and bottom to top of the periodic table.
 - the attraction of the nucleus to the outermost electron remains constant for each element on the periodic table.
22. Lithium's first and second ionization energies are 519 kJ/mol and 7300 kJ/mol, respectively. Element X has a first ionization energy of 590 kJ/mol and a second ionization energy of 1150 kJ/mol. Element X is most likely to be:
- Xenon
 - Fluorine
 - Calcium
 - Oxygen
 - Sodium
23. Which of the following is correctly ordered in terms of atomic radius, from smallest to largest?
- $F^- < Mg^{2+} < Na^+$
 - $Mg^{2+} < F^- < Na^+$
 - $F^- < Na^+ < Mg^{2+}$
 - $Mg^{2+} < Na^+ < F^-$
 - $Na^+ < F^- < Mg^{2+}$
24. Arrange the following in order of increasing electron affinity: Cl, P, S, Ar.
- Most positive EA $Ar < Cl < S < P$ Most negative EA
 - Most positive EA $Cl < S < P < Ar$ Most negative EA
 - Most positive EA $P < S < Cl < Ar$ Most negative EA
 - Most positive EA $Ar < P < S < Cl$ Most negative EA
 - Most positive EA $Cl < Ar < P < S$ Most negative EA
25. Which of the following statements is TRUE?
- The atomic radius of Li is larger than that of Cs.
 - The electron affinity of C is greater than that of Cl.
 - The ionic radius of Fe^+ is larger than that of Fe^{3+} .
 - The first ionization potential of H is greater than that of He.
 - All are false.

Version: C

1. Energy is released when nitrogen gas combines with hydrogen gas to form ammonia according to the equation:




Which of the following is responsible for the release of energy?

- A. Forming nitrogen-hydrogen bonds
 - B. Breaking hydrogen bonds
 - C. Breaking nitrogen bonds
 - D. Breaking both nitrogen and hydrogen bonds.
 - E. All of these release energy
2. The standard enthalpy of formation of atomic oxygen is:
- A. Zero, since oxygen is an element and Positive, since breaking of bonds is always endothermic
 - B. Positive, since breaking of bonds is always endothermic
 - C. Zero, since oxygen is an element
 - D. Negative, zero, or positive depending on the temperature
 - E. Negative, since all spontaneous reactions are exothermic
3. Given the following standard enthalpies of formation:

Compound	ΔH_f (kJ/mol)
$\text{CH}_4(\text{g})$	-75
$\text{CO}_2(\text{g})$	-394
$\text{H}_2\text{O}(\text{g})$	-242


What is the standard enthalpy change for the combustion of methane:




- A. -561 kJ
- B. +561 kJ
- C. +803 kJ
- D. -711 kJ
- E. -803 kJ

4. The standard enthalpy of formation for $\text{H}_2\text{O}(\text{l})$ is $\Delta\text{H} = -285 \text{ kJ/mol}$.

Beginning with $\text{H}_2\text{O}(\text{l})$, which of the following physical and/or chemical changes is/are **exothermic**?



- A. i, ii & iii
 B. only ii
 C. ii & iv
 D. only i
 E. only iv
5. Which of the following are **NOT** correctly written as a standard enthalpy of formation reactions?




- A. iii & iv
 B. i & iv
 C. i & ii
 D. ii, iii & iv
 E. all are correct

6. 100 mL of water, 100 mL of alcohol, and 200 mL of an unknown liquid, all at 25°C, are heated at the **same rate under identical conditions**. After 3 minutes the temperature of the alcohol is 50°C. It took 5 minutes for the water to reach 50°C. It took 6 minutes for the unknown liquid to reach 50°C.

Which statement is TRUE regarding the specific heat, C_s ?

- A. C_s (water) > C_s (alcohol)
- B. C_s (alcohol) > C_s (unknown)
- C. C_s (water) < C_s (unknown)
- D. C_s (water) = C_s (alcohol)

7. What amount of heat energy, kJ, is released in the combustion of 16.0 g of C_3H_6 ?





- A. 504
- B. 624
- C. 725
- D. 581
- E. 783


8. Which of these will lead to the greatest increase in temperature when added to 100.0 mL of water at 25°C?

- A. 66 g of silver at 124°C, specific heat = 0.234 J/g K.
- B. 50 g of iron at 110°C, specific heat = 0.452 J/g K.
- C. 94 g of gold at 150°C, specific heat = 0.128 J/g K.
- D. 70 g of copper at 150°C, specific heat = 0.385 J/g K.

9. Given the following equations and ΔH° values, determine the heat of reaction (kJ) at 298 K for the reaction:



- A. -300.1
 B. +770.8
 C. -605.0
 D. +291.2
 E. -291.6
10. A volume of 50.0 mL of 0.400 M NaOH(aq) was added to 20.0 mL of 0.500 M $\text{H}_2\text{SO}_4(\text{aq})$ in a calorimeter of heat capacity 39.0 JK^{-1} . The temperature of the resulting solution rose by 3.60°C . What is the total heat produced (in kJ) in this neutralization? Assume that the specific heat capacity of the neutralized solution is $4.18 \text{ JK}^{-1}\text{g}^{-1}$.



- A. -119
 B. -111
 C. -102
 D. -126
 E. -132

11. In the introduction to Chapter 6 I played several clips from the hit TV show “The Big Bang Theory” in class. In this episode, Dr. Sheldon Cooper is trying to visualize the behavior of electrons traveling through a graphene sheet. Which of the following statements is/are true regarding his findings?

i) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model with balls from a ball pit after he determined the proper size ratio with bigger carbon atoms.

ii) Just like Dalton, Thompson, and Bohr, it was relatively easy for Sheldon to develop an accurate model of the behavior of an electron.

iii) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model after recognizing the electron behaved as both a particle and a wave.

- A. i & ii
- B. ii only
- C. ii & iii
- D. i & iii
- E. iii only


12. What is the energy of the photons emitted from the He-Ne laser, wavelength = 655 nm?

- A. 4.34×10^{-40} J
- B. 1.01×10^{-27} J
- C. 3.03×10^{-19} J
- D. 2.27×10^{-19} J
- E. 5.29×10^{-18} J

13. Which emission line in the hydrogen spectrum occurs at highest frequency?

- A. $n = 2$ to $n = 4$
- B. $n = 3$ to $n = 1$
- C. $n = 4$ to $n = 2$
- D. $n = 1$ to $n = 3$
- E. $n = 5$ to $n = 7$

14.



Suppose a certain atom has four energy levels, and assume all transitions between levels are possible. Which statement is TRUE?

- A. The longest wavelength light is emitted for $n = 4 \rightarrow n = 2$
 - B. The lowest energy light is emitted for $n=3 \rightarrow n = 1$
 - C. The potential energy is greatest when the electron is at the $n=1$ level.
 - D. There are a total of 6 spectral emission lines.
 - E. There are a total of 4 spectral emission lines.
15. Sodium metal requires a photon with a minimum energy of 4.41×10^{-19} J to emit electrons. Suppose you have two lasers, one emitting light with a wavelength of 210 nm and the other emitting light with a wavelength of 335 nm. Each laser produces 10,000 photons/sec.

Which statement is true regarding the removal of electrons from the metal's surface?


- A. Only the laser with photons of 210 nm ejects electrons.
- B. Only the laser with photons of 335 nm ejects electrons.
- C. Both lasers eject electrons from the metal surface, but the weaker laser needs to operate for a longer period of time.
- D. Both lasers eject electrons from the metal surface at the same rate.
- E. Neither laser will eject electrons.

16. Which of the following statements best describes a $4d_{xz}$ atomic orbital?
- The $4d_{xz}$ orbital has more angular nodes than radial nodes.
 - The $4d_{xz}$ orbital has the same number of radial and angular nodes.
 - The $4d_{xz}$ orbital does not have any radial nodes.
 - The $4d_{xz}$ orbital does not have any angular nodes.
 - The $4d_{xz}$ orbital has more radial nodes than angular nodes.
17. Which of the following electron configurations is correct for mercury (Hg)?
- $[\text{Xe}] 6s^2 4f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 5d^{10}$
 - $[\text{Xe}] 4s^2 5f^{14} 6d^{10}$
 - $[\text{Kr}] 6s^2 6f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 5d^{10}$
18. How many electrons in a zinc atom have the quantum number $m_l = 0$?
- 30
 - 6
 - 10
 - 14
 - 8
19. Which response correctly fills in the blanks?
- The atom _____ has _____ core electrons, _____ valence electrons, and _____ unpaired electrons.
- chlorine, 12, 5, 2
 - fluorine, 2, 7, 5
 - neon, 2, 8, 6
 - nitrogen, 2, 5, 3
 - oxygen, 2, 6, 4
20. Which of the following sets of quantum numbers could describe one of the valence electrons in a sulfur atom in the ground state?
- $n=2 \quad l=1 \quad m_l = -1 \quad m_s = -\frac{1}{2}$
 - $n=1 \quad l=0 \quad m_l = 0 \quad m_s = +\frac{1}{2}$
 - $n=3 \quad l=1 \quad m_l = 0 \quad m_s = +\frac{1}{2}$
 - $n=3 \quad l=3 \quad m_l = -2 \quad m_s = +\frac{1}{2}$
 - $n=4 \quad l=0 \quad m_l = -1 \quad m_s = +\frac{1}{2}$

21. The attraction of the nucleus on the outermost electron in an atom tends to:
- decrease moving from right to left and top to bottom on the periodic table.
 - the attraction of the nucleus to the outermost electron remains constant for each element on the periodic table.
 - decrease moving from left to right and bottom to top on the periodic table.
 - decrease moving from right to left and bottom to top of the periodic table.
 - decrease moving from left to right and top to bottom on the periodic table.
22. Lithium's first and second ionization energies are 519 kJ/mol and 7300 kJ/mol, respectively. Element X has a first ionization energy of 590 kJ/mol and a second ionization energy of 1150 kJ/mol. Element X is most likely to be:
- Sodium
 - Calcium
 - Oxygen
 - Xenon
 - Fluorine
23. Which of the following is correctly ordered in terms of atomic radius, from smallest to largest?
- $\text{Mg}^{2+} < \text{F}^- < \text{Na}^+$
 - $\text{Mg}^{2+} < \text{Na}^+ < \text{F}^-$
 - $\text{F}^- < \text{Na}^+ < \text{Mg}^{2+}$
 - $\text{F}^- < \text{Mg}^{2+} < \text{Na}^+$
 - $\text{Na}^+ < \text{F}^- < \text{Mg}^{2+}$
24. Arrange the following in order of increasing electron affinity: Cl, P, S, Ar.
- Most positive EA $\text{Ar} < \text{Cl} < \text{S} < \text{P}$ Most negative EA
 - Most positive EA $\text{P} < \text{S} < \text{Cl} < \text{Ar}$ Most negative EA
 - Most positive EA $\text{Cl} < \text{S} < \text{P} < \text{Ar}$ Most negative EA
 - Most positive EA $\text{Ar} < \text{P} < \text{S} < \text{Cl}$ Most negative EA
 - Most positive EA $\text{Cl} < \text{Ar} < \text{P} < \text{S}$ Most negative EA
25. Which of the following statements is TRUE?
- The electron affinity of C is greater than that of Cl.
 - The first ionization potential of H is greater than that of He.
 - The atomic radius of Li is larger than that of Cs.
 - The ionic radius of Fe^+ is larger than that of Fe^{3+} .
 - All are false.

Version: D

1. Energy is released when nitrogen gas combines with hydrogen gas to form ammonia according to the equation:




Which of the following is responsible for the release of energy?

- A. Breaking hydrogen bonds
 - B. Forming nitrogen-hydrogen bonds
 - C. Breaking nitrogen bonds
 - D. Breaking both nitrogen and hydrogen bonds.
 - E. All of these release energy
2. The standard enthalpy of formation of atomic oxygen is:
- A. Negative, zero, or positive depending on the temperature
 - B. Zero, since oxygen is an element
 - C. Negative, since all spontaneous reactions are exothermic
 - D. Zero, since oxygen is an element and Positive, since breaking of bonds is always endothermic
 - E. Positive, since breaking of bonds is always endothermic
3. Given the following standard enthalpies of formation:

Compound	ΔH_f (kJ/mol)
CH ₄ (g)	-75
CO ₂ (g)	-394
H ₂ O(g)	-242


What is the standard enthalpy change for the combustion of methane:




- A. -803 kJ
- B. -561 kJ
- C. +561 kJ
- D. +803 kJ
- E. -711 kJ

4. The standard enthalpy of formation for $\text{H}_2\text{O}(\text{l})$ is $\Delta\text{H} = -285 \text{ kJ/mol}$.

Beginning with $\text{H}_2\text{O}(\text{l})$, which of the following physical and/or chemical changes is/are **exothermic**?



- A. i, ii & iii
 B. ii & iv
 C. only ii
 D. only i
 E. only iv
5. Which of the following are **NOT** correctly written as a standard enthalpy of formation reactions?




- A. ii, iii & iv
 B. i & iv
 C. i & ii
 D. iii & iv
 E. all are correct

6. 100 mL of water, 100 mL of alcohol, and 200 mL of an unknown liquid, all at 25°C, are heated at the **same rate under identical conditions**. After 3 minutes the temperature of the alcohol is 50°C. It took 5 minutes for the water to reach 50°C. It took 6 minutes for the unknown liquid to reach 50°C.

Which statement is TRUE regarding the specific heat, C_s ?

- A. C_s (water) > C_s (alcohol)
- B. C_s (water) < C_s (unknown)
- C. C_s (water) = C_s (alcohol)
- D. C_s (alcohol) > C_s (unknown)

7. What amount of heat energy, kJ, is released in the combustion of 16.0 g of C_3H_6 ?





- A. 783
- B. 504
- C. 624
- D. 581
- E. 725


8. Which of these will lead to the greatest increase in temperature when added to 100.0 mL of water at 25°C?

- A. 94 g of gold at 150°C, specific heat = 0.128 J/g K.
- B. 70 g of copper at 150°C, specific heat = 0.385 J/g K.
- C. 50 g of iron at 110°C, specific heat = 0.452 J/g K.
- D. 66 g of silver at 124°C, specific heat = 0.234 J/g K.

9. Given the following equations and ΔH° values, determine the heat of reaction (kJ) at 298 K for the reaction:



- A. -605.0
 B. -300.1
 C. +770.8
 D. +291.2
 E. -291.6
10. A volume of 50.0 mL of 0.400 M NaOH(aq) was added to 20.0 mL of 0.500 M H_2SO_4 (aq) in a calorimeter of heat capacity 39.0 JK^{-1} . The temperature of the resulting solution rose by 3.60°C . What is the total heat produced (in kJ) in this neutralization? Assume that the specific heat capacity of the neutralized solution is $4.18 \text{ JK}^{-1}\text{g}^{-1}$.



- A. -111
 B. -119
 C. -126
 D. -102
 E. -132

11. In the introduction to Chapter 6 I played several clips from the hit TV show “The Big Bang Theory” in class. In this episode, Dr. Sheldon Cooper is trying to visualize the behavior of electrons traveling through a graphene sheet. Which of the following statements is/are true regarding his findings?

- i) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model with balls from a ball pit after he determined the proper size ratio with bigger carbon atoms.
- ii) Just like Dalton, Thompson, and Bohr, it was relatively easy for Sheldon to develop an accurate model of the behavior of an electron.
- iii) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model after recognizing the electron behaved as both a particle and a wave.

- A. iii only
- B. i & ii
- C. ii only
- D. i & iii
- E. ii & iii


12. What is the energy of the photons emitted from the He-Ne laser, wavelength = 655 nm?

- A. 4.34×10^{-40} J
- B. 1.01×10^{-27} J
- C. 2.27×10^{-19} J
- D. 3.03×10^{-19} J
- E. 5.29×10^{-18} J

13. Which emission line in the hydrogen spectrum occurs at highest frequency?

- A. $n = 1$ to $n = 3$
- B. $n = 3$ to $n = 1$
- C. $n = 4$ to $n = 2$
- D. $n = 2$ to $n = 4$
- E. $n = 5$ to $n = 7$

14.



Suppose a certain atom has four energy levels, and assume all transitions between levels are possible. Which statement is TRUE?

- A. The potential energy is greatest when the electron is at the $n=1$ level.
 - B. The lowest energy light is emitted for $n=3 \rightarrow n=1$
 - C. The longest wavelength light is emitted for $n=4 \rightarrow n=2$
 - D. There are a total of 6 spectral emission lines.
 - E. There are a total of 4 spectral emission lines.
15. Sodium metal requires a photon with a minimum energy of 4.41×10^{-19} J to emit electrons. Suppose you have two lasers, one emitting light with a wavelength of 210 nm and the other emitting light with a wavelength of 335 nm. Each laser produces 10,000 photons/sec.

Which statement is true regarding the removal of electrons from the metal's surface?


- A. Only the laser with photons of 335 nm ejects electrons.
- B. Neither laser will eject electrons.
- C. Only the laser with photons of 210 nm ejects electrons.
- D. Both lasers eject electrons from the metal surface at the same rate.
- E. Both lasers eject electrons from the metal surface, but the weaker laser needs to operate for a longer period of time.

16. Which of the following statements best describes a $4d_{xz}$ atomic orbital?
- The $4d_{xz}$ orbital has more angular nodes than radial nodes.
 - The $4d_{xz}$ orbital has more radial nodes than angular nodes.
 - The $4d_{xz}$ orbital has the same number of radial and angular nodes.
 - The $4d_{xz}$ orbital does not have any radial nodes.
 - The $4d_{xz}$ orbital does not have any angular nodes.
17. Which of the following electron configurations is correct for mercury (Hg)?
- $[\text{Xe}] 4s^2 5f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 6d^{10}$
 - $[\text{Kr}] 6s^2 6f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 5d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 5d^{10}$
18. How many electrons in a zinc atom have the quantum number $m_l = 0$?
- 10
 - 8
 - 6
 - 14
 - 30
19. Which response correctly fills in the blanks?
- The atom _____ has _____ core electrons, _____ valence electrons, and _____ unpaired electrons.
- neon, 2, 8, 6
 - chlorine, 12, 5, 2
 - nitrogen, 2, 5, 3
 - oxygen, 2, 6, 4
 - fluorine, 2, 7, 5
20. Which of the following sets of quantum numbers could describe one of the valence electrons in a sulfur atom in the ground state?
- $n=3 \quad l=3 \quad m_l = -2 \quad m_s = +\frac{1}{2}$
 - $n=3 \quad l=1 \quad m_l = 0 \quad m_s = +\frac{1}{2}$
 - $n=2 \quad l=1 \quad m_l = -1 \quad m_s = -\frac{1}{2}$
 - $n=1 \quad l=0 \quad m_l = 0 \quad m_s = +\frac{1}{2}$
 - $n=4 \quad l=0 \quad m_l = -1 \quad m_s = +\frac{1}{2}$

21. The attraction of the nucleus on the outermost electron in an atom tends to:
- decrease moving from right to left and top to bottom on the periodic table.
 - decrease moving from right to left and bottom to top of the periodic table.
 - decrease moving from left to right and bottom to top on the periodic table.
 - decrease moving from left to right and top to bottom on the periodic table.
 - the attraction of the nucleus to the outermost electron remains constant for each element on the periodic table.
22. Lithium's first and second ionization energies are 519 kJ/mol and 7300 kJ/mol, respectively. Element X has a first ionization energy of 590 kJ/mol and a second ionization energy of 1150 kJ/mol. Element X is most likely to be:
- Sodium
 - Calcium
 - Xenon
 - Oxygen
 - Fluorine
23. Which of the following is correctly ordered in terms of atomic radius, from smallest to largest?
- $F^- < Na^+ < Mg^{2+}$
 - $Mg^{2+} < F^- < Na^+$
 - $F^- < Mg^{2+} < Na^+$
 - $Mg^{2+} < Na^+ < F^-$
 - $Na^+ < F^- < Mg^{2+}$
24. Arrange the following in order of increasing electron affinity: Cl, P, S, Ar.
- Most positive EA $Ar < P < S < Cl$ Most negative EA
 - Most positive EA $Ar < Cl < S < P$ Most negative EA
 - Most positive EA $P < S < Cl < Ar$ Most negative EA
 - Most positive EA $Cl < Ar < P < S$ Most negative EA
 - Most positive EA $Cl < S < P < Ar$ Most negative EA
25. Which of the following statements is TRUE?
- The ionic radius of Fe^+ is larger than that of Fe^{3+} .
 - The atomic radius of Li is larger than that of Cs.
 - The first ionization potential of H is greater than that of He.
 - The electron affinity of C is greater than that of Cl.
 - All are false.

Version: E

1. Energy is released when nitrogen gas combines with hydrogen gas to form ammonia according to the equation:




Which of the following is responsible for the release of energy?

- A. Breaking hydrogen bonds
 - B. Breaking both nitrogen and hydrogen bonds.
 - C. All of these release energy
 - D. Forming nitrogen-hydrogen bonds
 - E. Breaking nitrogen bonds
2. The standard enthalpy of formation of atomic oxygen is:
- A. Zero, since oxygen is an element
 - B. Positive, since breaking of bonds is always endothermic
 - C. Negative, zero, or positive depending on the temperature
 - D. Negative, since all spontaneous reactions are exothermic
 - E. Zero, since oxygen is an element and Positive, since breaking of bonds is always endothermic
3. Given the following standard enthalpies of formation:

Compound	ΔH_f (kJ/mol)
$\text{CH}_4(\text{g})$	-75
$\text{CO}_2(\text{g})$	-394
$\text{H}_2\text{O}(\text{g})$	-242


What is the standard enthalpy change for the combustion of methane:




- A. +803 kJ
- B. +561 kJ
- C. -561 kJ
- D. -803 kJ
- E. -711 kJ

4. The standard enthalpy of formation for $\text{H}_2\text{O}(\text{l})$ is $\Delta\text{H} = -285 \text{ kJ/mol}$.

Beginning with $\text{H}_2\text{O}(\text{l})$, which of the following physical and/or chemical changes is/are **exothermic**?



- A. ii & iv
 B. only i
 C. only iv
 D. only ii
 E. i, ii & iii
5. Which of the following are **NOT** correctly written as a standard enthalpy of formation reactions?




- A. ii, iii & iv
 B. i & ii
 C. i & iv
 D. iii & iv
 E. all are correct

6. 100 mL of water, 100 mL of alcohol, and 200 mL of an unknown liquid, all at 25°C, are heated at the **same rate under identical conditions**. After 3 minutes the temperature of the alcohol is 50°C. It took 5 minutes for the water to reach 50°C. It took 6 minutes for the unknown liquid to reach 50°C.

Which statement is TRUE regarding the specific heat, C_s ?

- A. C_s (water) = C_s (alcohol)
- B. C_s (water) < C_s (unknown)
- C. C_s (water) > C_s (alcohol)
- D. C_s (alcohol) > C_s (unknown)

7. What amount of heat energy, kJ, is released in the combustion of 16.0 g of C_3H_6 ?





- A. 624
- B. 783
- C. 504
- D. 581
- E. 725


8. Which of these will lead to the greatest increase in temperature when added to 100.0 mL of water at 25°C?

- A. 70 g of copper at 150°C, specific heat = 0.385 J/g K.
- B. 50 g of iron at 110°C, specific heat = 0.452 J/g K.
- C. 94 g of gold at 150°C, specific heat = 0.128 J/g K.
- D. 66 g of silver at 124°C, specific heat = 0.234 J/g K.

9. Given the following equations and ΔH° values, determine the heat of reaction (kJ) at 298 K for the reaction:



- A. -605.0
 B. -291.6
 C. -300.1
 D. +291.2
 E. +770.8
10. A volume of 50.0 mL of 0.400 M NaOH(aq) was added to 20.0 mL of 0.500 M $\text{H}_2\text{SO}_4(\text{aq})$ in a calorimeter of heat capacity 39.0 J K^{-1} . The temperature of the resulting solution rose by 3.60°C . What is the total heat produced (in kJ) in this neutralization? Assume that the specific heat capacity of the neutralized solution is $4.18 \text{ J K}^{-1} \text{ g}^{-1}$.



- A. -126
 B. -119
 C. -102
 D. -111
 E. -132

11. In the introduction to Chapter 6 I played several clips from the hit TV show “The Big Bang Theory” in class. In this episode, Dr. Sheldon Cooper is trying to visualize the behavior of electrons traveling through a graphene sheet. Which of the following statements is/are true regarding his findings?

- i) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model with balls from a ball pit after he determined the proper size ratio with bigger carbon atoms.
- ii) Just like Dalton, Thompson, and Bohr, it was relatively easy for Sheldon to develop an accurate model of the behavior of an electron.
- iii) Sheldon’s model of the behavior of electrons failed with lima beans, peas, and marbles, but he was able to come up with an accurate model after recognizing the electron behaved as both a particle and a wave.

- A. iii only
- B. ii & iii
- C. i & ii
- D. i & iii
- E. ii only


12. What is the energy of the photons emitted from the He-Ne laser, wavelength = 655 nm?

- A. 3.03×10^{-19} J
- B. 4.34×10^{-40} J
- C. 5.29×10^{-18} J
- D. 2.27×10^{-19} J
- E. 1.01×10^{-27} J

13. Which emission line in the hydrogen spectrum occurs at highest frequency?

- A. $n = 5$ to $n = 7$
- B. $n = 4$ to $n = 2$
- C. $n = 2$ to $n = 4$
- D. $n = 1$ to $n = 3$
- E. $n = 3$ to $n = 1$

14.



Suppose a certain atom has four energy levels, and assume all transitions between levels are possible. Which statement is TRUE?

- A. The longest wavelength light is emitted for $n = 4 \rightarrow n = 2$
 - B. There are a total of 6 spectral emission lines.
 - C. The potential energy is greatest when the electron is at the $n=1$ level.
 - D. The lowest energy light is emitted for $n=3 \rightarrow n = 1$
 - E. There are a total of 4 spectral emission lines.
15. Sodium metal requires a photon with a minimum energy of 4.41×10^{-19} J to emit electrons. Suppose you have two lasers, one emitting light with a wavelength of 210 nm and the other emitting light with a wavelength of 335 nm. Each laser produces 10,000 photons/sec.

Which statement is true regarding the removal of electrons from the metal's surface?

- A. Both lasers eject electrons from the metal surface, but the weaker laser needs to operate for a longer period of time.
- B. Both lasers eject electrons from the metal surface at the same rate.
- C. Only the laser with photons of 335 nm ejects electrons.
- D. Only the laser with photons of 210 nm ejects electrons.
- E. Neither laser will eject electrons.

16. Which of the following statements best describes a $4d_{xz}$ atomic orbital?
- The $4d_{xz}$ orbital has more radial nodes than angular nodes.
 - The $4d_{xz}$ orbital has more angular nodes than radial nodes.
 - The $4d_{xz}$ orbital has the same number of radial and angular nodes.
 - The $4d_{xz}$ orbital does not have any radial nodes.
 - The $4d_{xz}$ orbital does not have any angular nodes.
17. Which of the following electron configurations is correct for mercury (Hg)?
- $[\text{Xe}] 4s^2 5f^{14} 6d^{10}$
 - $[\text{Kr}] 6s^2 6f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 6d^{10}$
 - $[\text{Xe}] 6s^2 5d^{10}$
 - $[\text{Xe}] 6s^2 4f^{14} 5d^{10}$
18. How many electrons in a zinc atom have the quantum number $m_l = 0$?
- 10
 - 14
 - 6
 - 8
 - 30
19. Which response correctly fills in the blanks?
- The atom _____ has _____ core electrons, _____ valence electrons, and _____ unpaired electrons.
- nitrogen, 2, 5, 3
 - neon, 2, 8, 6
 - oxygen, 2, 6, 4
 - chlorine, 12, 5, 2
 - fluorine, 2, 7, 5
20. Which of the following sets of quantum numbers could describe one of the valence electrons in a sulfur atom in the ground state?
- $n=3 \quad l=1 \quad m_l=0 \quad m_s=+\frac{1}{2}$
 - $n=4 \quad l=0 \quad m_l=-1 \quad m_s=+\frac{1}{2}$
 - $n=2 \quad l=1 \quad m_l=-1 \quad m_s=-\frac{1}{2}$
 - $n=3 \quad l=3 \quad m_l=-2 \quad m_s=+\frac{1}{2}$
 - $n=1 \quad l=0 \quad m_l=0 \quad m_s=+\frac{1}{2}$

21. The attraction of the nucleus on the outermost electron in an atom tends to:
- decrease moving from left to right and bottom to top on the periodic table.
 - decrease moving from right to left and bottom to top of the periodic table.
 - decrease moving from left to right and top to bottom on the periodic table.
 - decrease moving from right to left and top to bottom on the periodic table.
 - the attraction of the nucleus to the outermost electron remains constant for each element on the periodic table.
22. Lithium's first and second ionization energies are 519 kJ/mol and 7300 kJ/mol, respectively. Element X has a first ionization energy of 590 kJ/mol and a second ionization energy of 1150 kJ/mol. Element X is most likely to be:
- Calcium
 - Sodium
 - Oxygen
 - Xenon
 - Fluorine
23. Which of the following is correctly ordered in terms of atomic radius, from smallest to largest?
- $F^- < Mg^{2+} < Na^+$
 - $F^- < Na^+ < Mg^{2+}$
 - $Mg^{2+} < F^- < Na^+$
 - $Mg^{2+} < Na^+ < F^-$
 - $Na^+ < F^- < Mg^{2+}$
24. Arrange the following in order of increasing electron affinity: Cl, P, S, Ar.
- Most positive EA $Ar < Cl < S < P$ Most negative EA
 - Most positive EA $Cl < S < P < Ar$ Most negative EA
 - Most positive EA $P < S < Cl < Ar$ Most negative EA
 - Most positive EA $Ar < P < S < Cl$ Most negative EA
 - Most positive EA $Cl < Ar < P < S$ Most negative EA
25. Which of the following statements is TRUE?
- The electron affinity of C is greater than that of Cl.
 - The first ionization potential of H is greater than that of He.
 - The atomic radius of Li is larger than that of Cs.
 - The ionic radius of Fe^+ is larger than that of Fe^{3+} .
 - All are false.